



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# CONCERNING APPROACHABILITY OF SIMPLE CLOSED AND OPEN CURVES\*

BY

JOHN ROBERT KLINE

Schoenflies† was the first to formulate the converse of the fundamental theorem of C. Jordan‡ that a simple closed curve§ lying wholly within a plane decomposes the plane into an inside and an outside region. The statement of this converse theorem is as follows: *Suppose  $K$  is a closed, bounded set of points lying in a plane  $S$  and that  $S - K = S_1 + S_2$ , where  $S_1$  and  $S_2$  are point-sets such that (1) every two points of  $S_i$  ( $i = 1, 2$ ) can be joined by an arc lying entirely in  $S_i$  (2) every arc joining a point of  $S_1$  to a point of  $S_2$  contains at least one point of  $K$  (3) if  $O$  is a point of  $K$  and  $P$  is a point not belonging to  $K$ , then  $P$  can be joined to  $O$  by an arc that has no point except  $O$  in common with  $K$ . Every point-set that satisfies these conditions is a simple closed curve.* Schoenflies used metrical hypotheses in his proof. Lennes gave a proof of this converse theorem using straight lines.|| R. L. Moore pointed out that a proof similar in large part to that of Lennes can be carried through with the use of arcs and closed curves on the basis of his system of postulates  $\Sigma_3$ , thus furnishing a non-metrical proof of the converse theorem.¶

In all these proofs the condition numbered three, the condition of approachability (*erreichbarkeit*) plays a fundamental rôle. It is the purpose of the present paper to study the effect of substituting for the condition of approach-

\* Presented to the Society, April, 1918.

† Cf. A. Schoenflies, *Ueber einen grundlegenden Satz der Analysis Situs*, *Nachrichten der Göttinger Gesellschaft der Wissenschaften*, 1902, p. 185.

‡ C. Jordan, *Cours d'Analyse*, 2d ed., Paris, 1893, p. 92.

§ If  $A$  and  $B$  are distinct points, then a *simple continuous arc* from  $A$  to  $B$  is defined by Lennes as a bounded, closed, connected set of points containing  $A$  and  $B$ , but containing no proper connected subset containing both  $A$  and  $B$ . Cf. N. J. Lennes *Curves in non-metrical analysis situs with an application in the calculus of variations*, *American Journal of Mathematics*, vol. 33 (1911), p. 308. A *simple closed curve* is a set of points composed of two arcs  $AXB$  and  $AYB$  having no point in common other than  $A$  and  $B$ . Hereafter in this paper "arc" and "closed curve" will be considered synonymous with "simple continuous arc" and "simple closed curve," respectively.

|| Cf. N. J. Lennes, loc. cit., § 5.

¶ Cf. R. L. Moore, *On the foundations of plane analysis situs*, these *Transactions*, vol. 17 (1916), p. 59.

ability, the condition that the set is "*connected in kleinem*."\* The results obtained are embodied in the following theorem:

**THEOREM A.** *Suppose  $K$  is a closed plane point-set,  $S$  is the set of all points of the plane, while  $S - K = S_1 + S_2$ , where  $S_1$  and  $S_2$  are two mutually exclusive domains† such that every point of  $K$  is a common boundary point of  $S_1$  and  $S_2$ . Then a necessary and sufficient condition that  $K$  be either a simple closed curve or an open curve‡ is that  $K$  be connected in kleinem.*

That the condition stated in Theorem A is necessary is evident. I will proceed to show that it is sufficient. Suppose  $K$  is a connected in kleinem set satisfying the conditions stipulated in Theorem A. Then the following lemmas hold true:

**LEMMA A.** *Every arc joining a point of  $S_1$  to a point of  $S_2$  contains a point of  $K$ .*

*Proof.* Suppose it were possible to draw an arc from a point  $P_1$  of  $S_1$  to a point  $P_2$  of  $S_2$  that contains no point of  $K$ . Then let us divide the arc  $P_1 P_2$  into two sets,  $M_1$  and  $M_2$ , where  $M_1$  is the set of all points of  $P_1 P_2$  that belong to  $S_1$ , while  $M_2$  is the set of all points of  $P_1 P_2$  which belong to  $S_2$ . As  $P_1 P_2$  is a connected point-set either  $M_1$  contains a limit point of  $M_2$  or  $M_2$  contains a limit point of  $M_1$ .

*Case I.* A point  $F$  of  $M_1$  is a limit point of  $M_2$ . As  $F$  is a point of the domain  $S_1$ , there exists a region containing  $F$  and lying entirely in  $S_1$ . As  $S_1$  and  $S_2$  are mutually exclusive domains, this region contains no point of  $M_2$ . Hence  $F$  cannot be a limit point of  $M_2$ .

*Case II.* A point  $G$  of  $M_2$  is a limit point of  $M_1$ . This is impossible as in Case I.

Hence we are led to a contradiction if we suppose our lemma false.

**LEMMA B.** *The set  $K$  is connected.*

*Proof.* Suppose  $K$  were not connected. Then it could be divided into two mutually exclusive sets  $K_1$  and  $K_2$ , neither of which contains a limit point of the other one. Let  $P_i$  ( $i = 1, 2$ ) denote a point of  $K_i$ . Put about  $P_i$  a circle  $R_i$  having  $P_i$  as center and such that  $R_i$  and its interior lie entirely

\* Cf. Hans Hahn, *Ueber die allgemeinste ebene Punktmenge die stetiges Bild einer Strecke ist*, Jahresbericht der Deutschen Mathematiker Vereinigung, vol. 23 (1914), pp. 318-22. According to Hahn, a set of points  $C$  is said to be *connected in kleinem* if, whenever  $P$  is a point of  $C$ ,  $\epsilon$  a positive number and  $K$  a circle of radius  $1/\epsilon$  with center at  $P$ , then there exists within  $K$  and with center at  $P$ , another circle  $K_{\epsilon, P}$  such that if  $X$  is a point of  $C$  within  $K_{\epsilon, P}$  then  $X$  and  $P$  lie together in some connected subset of  $C$  that lies entirely within  $K$ .

† A *domain* is a connected set of points  $M$  such that if  $P$  is a point of  $M$ , then there is a region that contains  $P$  and lies in  $M$ .

‡ An *open curve* is defined by R. L. Moore as a closed, connected, set of points  $M$  such that if  $P$  is a point of  $M$ , then  $M - P$  is the sum of two mutually exclusive connected point-sets, neither of which contains a limit point of the other.

without  $R_{i+1}$ .<sup>\*</sup> As  $K$  is connected in kleinem, there exists a circle  $\bar{R}$ , lying within  $R_i$  and with center at  $P_i$  such that if  $X_i$  is a point of  $K$  within  $\bar{R}_i$ , then  $X_i$  and  $P_i$  lie on some connected subset of  $K$  lying within  $R_i$ . It may easily be shown that  $X_i$  can be joined to  $P_i$  by a simple continuous arc of  $K$  lying entirely within  $R_i$ .<sup>†</sup> As every point of  $K$  is a common boundary point of  $S_1$  and  $S_2$ , then there exists within  $\bar{R}_i$  a point  $M_{ij}$  ( $j = 1, 2$ ) belonging to  $S_j$ . As  $S_j$  is a domain, then there exists a simple continuous arc  $M_{1j} K_j M_{2j}$  lying entirely in  $S_j$ . Join  $M_{ij}$  to  $P_i$  by a simple continuous arc  $M_{ij} L_{ij} P_i$  lying entirely within  $\bar{R}_i$  and let  $G_{ij}$  denote the first point of  $K$  on the arc  $M_{ij} L_{ij} P_i$  following  $M_{ij}$ . Then we may join  $G_{i1}$  to  $G_{i2}$  by an arc  $G_{i1} F_i G_{i2}$  belonging to  $K$  and lying entirely within  $R_i$ . The point-set  $G_{11} M_{11}$  (on  $M_{11} L_{11} P_1$ ) +  $M_{11} K_1 M_{21}$  +  $M_{21} G_{21}$  (on  $M_{21} L_{21} P_2$ ) contains as a subset a simple continuous arc  $G_{11} H_1 G_{21}$  lying except for its endpoints entirely in  $S_1$ , while the set  $G_{12} M_{12}$  (on  $M_{12} L_{12} P_1$ ) +  $M_{12} K_2 M_{22}$  +  $M_{22} G_{22}$  (on  $M_{22} L_{22} P_2$ ) contains as a subset a simple continuous arc  $G_{12} H_2 G_{22}$  lying except for its endpoints entirely in  $S_2$ . We then have a closed curve  $G_{11} F_1 G_{12} H_2 G_{22} - F_2 G_{21} H_1 G_{11}$  such that the arcs  $G_{11} F_1 G_{12}$  and  $G_{21} F_2 G_{22}$  lie entirely on  $K$  and within  $R_1$  and  $R_2$ , respectively, while  $G_{11} H_1 G_{21}$ <sup>‡</sup> and  $G_{12} H_2 G_{22}$  belong to  $S_1$  and  $S_2$ , respectively.

All points of  $G_{11} F_1 G_{12}$  belong to  $K_1$ . For suppose a point  $H$  of  $G_{11} F_1 G_{12}$  belonged to  $K_2$ . As  $H$  is joined to  $G_{11}$ , which in turn can be joined to  $P_1$  by an arc of  $K$  lying entirely within  $R_1$ , it follows that  $H$  can be joined to  $P_1$  by an arc  $HFP_1$  of  $K$  lying entirely within  $R_1$ . Let  $[\bar{H}_1]$  denote the set of all points of  $HFP_1$  belonging to  $K_1$  while  $[\bar{H}_2]$  denotes the set of all points of  $HFP_1$  belonging to  $K_2$ . Clearly neither of these sets contains a limit point of the other. Hence the arc  $HFP_1$  is not a connected point-set. Hence the supposition that  $H$  belongs to  $K_2$  has led to a contradiction. In like manner, all points of  $G_{21} F_2 G_{22}$  belong to  $K_2$ .

The interior of  $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$  must contain at least one point of  $K$ . For suppose it does not contain a point of  $K$ . Then the interior of  $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$  is a subset of  $S_1 + S_2$ . Suppose it contains a point  $H$  of  $S_1$ . Then  $H$  can be joined to  $H_2$  by an arc  $HXH_2$  lying except for  $H_2$  entirely within  $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$ .<sup>§</sup> Let  $[W_1]$  denote the set of all points of  $HXH_2$  belonging to  $S_1$  while  $[W_2]$  denotes the set of all points of  $HXH_2$  which are points of  $S_2$ . Clearly neither of these sets contains

<sup>\*</sup> It is understood that subscripts are reduced modulo 2,

<sup>†</sup> Cf. R. L. Moore, *A theorem concerning continuous curves*, Bulletin of the American Mathematical Society, vol. 23 (1917). While Professor Moore's theorem states that every two points of a continuous curve can be joined by a simple continuous arc lying entirely on the given continuous curve, it is clear that his methods suffice to prove the above stronger statement.

<sup>‡</sup> If  $AXB$  is an arc, then the symbol  $AXB$  will denote  $AXB - A - B$ ,

<sup>§</sup> Cf. R. L. Moore, *Foundations of plane analysis situs*, loc. cit., Theorem 39, pp. 153-5.

a limit point of the other. Hence the arc  $HXH_2$  is not a connected point-set. In like manner the supposition that there is within  $G_{11} F_1 G_{12} H_2 G_{22} F_2 - G_{21} H_1 G_{11}$ , a point of  $S_2$  leads to a contradiction. Hence  $G_{11} F_1 G_{12} H_2 G_{22} F_2 - G_{21} H_1 G_{11}$  must enclose a point of  $K$ .

Let  $[V_2]$  denote the set of all points  $V_2$  such that either (1)  $V_2$  is a point of  $G_{21} F_2 G_{22}$ , or (2)  $V_2$  is a point such that there exists a closed connected set  $V_2 XF'_2$  belonging to  $K$  and lying within or on  $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$  and such that  $F'_2$  is a point of  $G_{21} F_2 G_{22}$ . As  $K$  is connected in kleinem it may easily be proved that  $[V_2]$  is a closed set. It is also true that all points of  $[V_2]$  belong to  $K_2$ . Hence no point of  $G_{11} F_1 G_{12}$  either belongs to or is a limit point of  $[V_2]$ . It may also be proved with the use of the in kleinem property that no point of  $[V_2]$  is a limit point of a set of points of  $K$  lying within  $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$  and containing no point of  $V_2$ . There exists an arc  $H_1 YH_2$  such that (1)  $H_1 YH_2$  is a subset of the interior of  $G_{11} F_1 G_{12} H_2 G_{22} F_2 G_{21} H_1 G_{11}$  and (2)  $\overline{H_1 YH_2}$  contains no points of  $[V_2]$ .<sup>\*</sup> Let  $[V_1]$  denote the set of all points of  $K$  within or on the closed curve,  $H_1 YH_2 G_{22} F_2 G_{21} H_1$ , not belonging to  $[V_2]$ . The set  $[V_1]$  is closed. Put about each point of  $[V_1]$  a circle lying entirely within  $G_{11} F_1 G_{12} H_2 G_{22} F_2 - G_{21} H_1 G_{11}$  and containing within it or on its boundary no point of  $[V_2]$ . By the Heine-Borel Property, there exists a finite number of circles of the above set,  $C_1, C_2, \dots, C_n$ , covering  $[V_1]$ . With the use of Theorems 41, 42, 43, and 44 of Professor Moore's Foundations we may easily obtain from the set  $C_1, C_2, \dots, C_n$  and the closed curve  $G_{11} F_1 G_{12} H_2 YH_1 G_{11}$ , a new closed curve  $G_{11} F_1 G_{12} H_2 ZH_1 G_{11}$ , where the arc  $H_1 G_{11} F_1 G_{12} H_2$  of the new closed curve  $G_{11} F_1 G_{12} H_2 ZH_1 G_{11}$  is the arc  $H_1 G_{11} F_1 G_{12} H_2$  of  $G_{11} F_1 G_{12} H_2 YH_1 G_{11}$  and where  $\overline{H_2 ZH_1}$  is free from points of  $K$  and lies within  $G_{11} F_1 G_{12} H_2 G_{22} F_2 - G_{21} H_1 G_{11}$ . But then we have a point of  $S_1$  joined to a point of  $S_2$  by an arc containing no point of  $K$ . Thus the supposition that  $K$  is not connected, leads to a contradiction.

LEMMA C. *If  $K$  contains one simple closed curve  $J$ , then all points of  $K$  belong to  $J$ .*

*Proof.* Suppose Lemma C is not true. Then  $K$  contains a closed curve  $J$  and at least one point  $P$  not on  $J$ . Two cases may arise:

*Case I.*  $P$  is within  $J$ . As every point of  $K$  is a common boundary point of  $S_1$  and  $S_2$ , the interior of  $J$  contains a point  $P_1$  of  $S_1$  and a point  $P_2$  of  $S_2$ . The exterior of  $J$  cannot contain a point  $\bar{P}_1$  of  $S_1$ . For suppose it did. Then any arc from  $P_1$  to  $\bar{P}_1$  would contain a point of  $J$  and hence a point of  $K$ , contrary to the fact that  $S_1$  is a domain. In like manner no point of  $S_2$  can be in the exterior of  $J$ . Hence the exterior of  $J$  must be a subset of  $K$ , while

<sup>\*</sup> Cf. my paper, *A definition of sense on plane closed curves in non-metrical analysis situs*, *Annals of Mathematics*, vol. XIX (1918), Theorem D, pp. 188-9.

$S_1$  and  $S_2$  are subsets of the interior of  $J$ . But this is impossible because no point without  $J$  is a limit point of a set of points lying entirely within  $J$  thus making it impossible that every point of  $K$  be a common boundary point of  $S_1$  and  $S_2$ . Hence the supposition that  $P$  is within  $J$  has led to a contradiction.

*Case II.*  $P$  is without  $J$ . Case II may be proved impossible by an argument similar to that used in Case I.

An immediate consequence of Lemma *C* is that if  $K$  is not a simple closed curve, then there is but one  $K$ -arc from a point  $A$  of  $K$  to a distinct point  $B$  of  $K$ .

LEMMA *D.* *The set  $K$  does not contain three arcs  $OP_1$ ,  $OP_2$ , and  $OP_3$ , no two of which have a common point other than  $O$ .*

*Proof.* Suppose Lemma *D* were false. Then there would exist three arcs  $OP_1$ ,  $OP_2$ , and  $OP_3$ , no two of which have a point in common other than  $O$ . Put about  $P_i$  ( $i = 1, 2, 3$ ) a circle  $C_i$  such that the point-set  $OP_{i+1}^* + OP_{i+2}$  is a subset of the exterior of  $C_i$  and such that  $C_i$  has no point in common with  $C_{i+1} + C_{i+2}$ . As  $K$  is connected in kleinem, there exists within  $C_i$  and with center at  $P_i$ , another circle  $C_{P_i, q_i}$  such that if  $X_i$  is a point of  $K$  within  $C_{P_i, q_i}$ , then there is an arc from  $X_i$  to  $P_i$  every point of which is a point of  $K$  and which lies entirely within  $C_i$ .† As all points of  $K$  are limit points of both  $S_1$  and  $S_2$ ,  $C_{P_i, q_i}$  must contain at least one point  $P_{i, 1}$  of  $S_1$ . As  $S_1$  is a domain, there is an arc  $P_{11}P_{21}$  from  $P_{11}$  to  $P_{21}$  all points of which belong to  $S_1$ . Join  $P_{i, 1}$  to  $P_i$  by an arc  $P_{i, 1}P_i$  lying entirely within  $C_{P_i, q_i}$  and let  $X_i$  denote the first point of the arc  $P_{i, 1}P_i$  after  $P_{i, 1}$ , which belongs to  $K$ . There exists an arc  $X_iP_i$  from  $X_i$  to  $P_i$  belonging to  $K$  and lying entirely within  $C_i$ . Let  $P'_i$  denote the first point of the arc  $X_iP_i$  which is on  $OP_i$ . The point-set  $P'_1X_1 + X_1P_{11} + P_{11}P_{21} + P_{21}X_2 + X_2P'_2$  contains as a subset an arc  $P'_1F_1P'_2$  such that (1)  $\overline{P'_1F_1P'_2}$  has no point in common with  $OP_1 + OP_2 + OP_3$ , (2) all points of  $\overline{P'_1F_1P'_2}$  belong to either  $K$  or  $S_1$ , (3) at least one point,  $F_1$ , of  $S_1$  is a point of  $P'_1F_1P'_2$ . By methods similar to those just employed, we may construct an arc  $Q'_1H_2Q'_2$  from a point  $Q'_1$  of  $OP_1$  to a point  $Q'_2$  of  $OP'_2$  such that (1)  $Q'_i$  is on  $OP_i$  between  $P'_i$  and  $O$ , (2) all points of  $Q'_1H_2Q'_2$  belong to either  $S_2$  or  $K$ , (3) except for  $Q'_1$  and  $Q'_2$ ,  $Q'_1H_2Q'_2$  has no point in common with  $P'_1FP'_2 + OP_1 + OP_2 + OP_3$ , (4) at least one point  $H_2$  of  $Q'_1H_2Q'_2$  belongs to  $S_2$ . Two cases may arise:

*Case I.*  $Q'_1H_2Q'_2$  is entirely within  $OP'_1F_1P'_2O$ . Then the interior of  $OP'_1F_1P'_2O = \overline{Q'_1H_2Q'_2} +$  the interior of  $OQ'_1H_2Q'_2O +$  the interior of  $P'_1F_1P'_2Q'_2H_2Q'_1P'_1$ . The point-set  $\overline{OP_3} + P_3$  is either entirely within or entirely without  $\overline{OQ'_1H_2Q'_2O}$ .

(a) Suppose  $\overline{OP_3} + P_3$  is entirely within  $\overline{OQ'_1H_2Q'_2O}$ . Then  $\overline{OQ'_1H_2Q'_2O}$

\* It is understood throughout this argument that subscripts are reduced modulo 3.

† See an earlier footnote.

must enclose at least one point  $L$  of  $S_1$ . But then an arc from  $L$  to  $F_1$  must contain at least one point of  $OQ'_1 H_2 Q'_2 O$ . Hence, as  $OQ'_1 H_2 Q'_2 O$  is a subset of  $K + S_2$ , no such arc  $LF_1$  can lie entirely in  $S_1$ , contrary to the fact that  $S_1$  is a domain.

(b) Suppose  $OP_3 + P_3$  is entirely without  $OQ'_1 H_2 Q'_2 O$ . It follows that  $OP_3 + P_3$  is entirely without  $OP'_1 F_1 P'_2 O$ . Then the exterior of  $OP'_1 F_1 P'_2 O$  contains at least one point  $M$  of  $S_2$ . Then any arc from  $M$  to  $H_2$  must contain at least one point of  $OP'_1 F_1 P'_2 O$  and hence at least one point not in  $S_2$ . But this is contrary to the fact that  $S_2$  is a domain.

Thus in Case I we are led to a contradiction.

*Case II.*  $Q'_1 H_2 Q'_2$  is without  $OP'_1 F_1 P'_2 O$ . We may show that Case II is impossible by methods similar to those used in Case I.

**LEMMA E.** *If  $O$  is a point of  $K$  and  $P$  is a point of  $S_i$  ( $i = 1, 2$ ) then there exists at least one arc  $OP$  such that  $\overline{OP} + P$  is a subset of  $S_i$ .*

*Proof.* Two conceivable cases may arise.

*Case I.* There exist points  $A_1$  and  $A_2$  of  $K \setminus [A_1 \neq O \neq A_2]$  such that  $O$  is a point of the arc  $A_1 OA_2$  belonging to  $K$ . By the same methods as were used in the preceding lemma we may construct an arc  $A'_1 F_1 A'_2$  such that (1) on  $A_1 OA_2$  the order  $A_1 A'_1 OA'_2 A_2$  holds, (2)  $A'_1 F_1 A'_2$  is a subset of  $S_1 + K$ , (3) at least one point  $F_1$  of  $A'_1 F_1 A'_2$  is a point of  $S_1$ , (4) no point of  $A'_1 F_1 A'_2$  belongs to  $A_1 OA_2$ . The point  $O$  is not a limit point of  $K - A'_1 OA'_2$ . For suppose it were. Then it would be a sequential limit point of a set of points  $P_1, P_2, \dots$ , every one of which belongs to  $K - A'_1 OA'_2$ . Put about  $O$  as center a circle  $M$  such that  $A'_1$  and  $A'_2$  are both without  $M$ . As  $K$  is connected in kleinem there exists another circle  $\bar{M}$  lying within  $M$  and having its center at  $O$  such that if  $X$  is a point of  $K$  within  $\bar{M}$ , then  $X$  and  $O$  can be joined by an arc of  $K$  lying entirely within  $M$ . Let  $\bar{P}$  denote that point of the set  $P_1, P_2, \dots$  of lowest subscript which lies within  $\bar{M}$ , while  $\bar{PO}$  denotes an arc of  $K$  from  $\bar{P}$  to  $O$  lying entirely within  $M$ . Let  $O'$  denote the first point of  $\bar{PO}$  which is on  $A'_1 OA'_2$ . Then the set  $K$  contains three arcs  $A'_1 O'$ ,  $A'_2 O'$ , and  $PO'$ , no two of which have a point in common other than  $O'$ . But this is contrary to Lemma D. Hence  $O$  cannot be a limit point of  $K - A'_1 OA'_2$ . There exists a closed curve  $G$  enclosing  $O$  but enclosing no points of  $A'_1 F_1 A'_2 + [K - A'_1 OA'_2]$ . Then there exist two closed curves  $J'_1$  and  $J'_2$  such that (1) every point of  $J'_1$  or  $J'_2$  belongs either to  $G$  or to  $A'_1 F_1 A'_2 OA'_1$  (2)  $O$  is on  $J'_1$  and on  $J'_2$  (3) every point within  $J'_1$  is within  $A'_1 F_1 A'_2 OA'_1$  while every point within  $J'_2$  is without  $A'_1 F_1 A'_2 OA'_1$  (4) every point within either  $J'_1$  or  $J'_2$  is within  $G$ .\* It is clear that either the interior of  $J'_1$  or the interior of  $J'_2$  is a subset of  $S_1$  while the interior of the other of these two closed curves is a subset of  $S_2$ . Let  $J_1$  denote that one whose interior is a subset of  $S_1$  while

\* Cf. R. L. Moore, *Foundations*, Theorem 43, pp. 156-7.

$J_2$  denotes the one whose interior is a subset of  $S_2$ . Let  $E$  denote a point within  $J_1$ , while  $P_1$  is any other point of  $S_1$ . There exists an arc  $EO$  such that  $EO - O$  is a subset of the interior of  $J_1$ .<sup>\*</sup> As  $S_1$  is a domain, there is an arc  $EP_1$  lying entirely in  $S_1$ . The point-set  $EO + EP_1$  contains as a subset an arc from  $P_1$  to  $O$  lying except for  $O$  entirely in  $S_1$ . In like manner we may show that any point  $P_2$  of  $S_2$  can be joined to  $O$  by an arc lying except for  $O$  entirely in  $S_2$ .

*Case II.* There do not exist two distinct points  $A_1$  and  $A_2$  of  $K$  such that  $O$  is on an arc of  $K$  from  $A_1$  to  $A_2$ . Let  $A$  denote a point of  $K$  different from  $O$  while  $ARO$  denotes an arc of  $K$  from  $A$  to  $O$ . By an argument similar to that employed in Case I we may show that if  $O$  were a limit point of  $K - ARO$ , then either there would exist three arcs  $AR'$ ,  $R'O$ , and  $R'P$ , no two of which have a point in common other than  $R'$  or there would exist a point  $A'$ , ( $A \neq A' \neq O$ ) such that  $O$  is an arc of  $K$  from  $A'$  to  $A$ . But the first of these possibilities contradicts Lemma *D* while the second is contrary to the hypothesis of Case II. Hence  $O$  cannot be a limit point of  $K - ARO$ . Put about  $O$  a circle  $C$  that neither contains or encloses any point of  $K - ARO$ . Let  $P_1, P_2, \dots$  denote a set of points of  $S_1$  approaching  $O$  as their sequential limit point. It is possible to pass at least one simple continuous arc<sup>†</sup> through  $ARO + P_1 + P_2 + \dots$ . Let  $P'_1ORA$  denote one such arc. If the interval  $OP'_1$  of the arc  $P'_1ORA$  does not lie entirely within  $C$ , let  $P'$  denote the first point which it has in common with  $C$ . Otherwise let  $P'$  denote  $P'_1$ . Let  $\overline{P}$  denote that point of the set  $P_1, P_2, \dots$  of lowest subscript lying on  $OP'$ . It is clear that the sub-arc  $OP'$  of  $P'_1ORA$  lies, except for  $O$ , entirely in  $\overline{S_1}$ . Let  $F_1$  denote any other point of  $S_1$ . Join  $F_1$  to  $\overline{P}$  by an arc lying entirely in  $S_1$ . Then the point-set  $OP' + \overline{P}F_1$  contains as a subset an arc from  $O$  to  $F_1$ , lying except for  $O$  entirely in  $S_1$ .

In like manner we may show that if  $F_2$  is a point of  $S_2$ ,  $F_2$  can be joined to  $O$  by an arc lying except for  $O$  entirely in  $S_2$ .

LEMMA *F*. *A necessary and sufficient condition that  $K$  be bounded, is that either  $S_1$  or  $S_2$  be bounded.*

*Proof.* *The condition is necessary.* Let us suppose that  $K$  is bounded while neither  $S_1$  nor  $S_2$  is bounded. As  $K$  is bounded, there is a circle  $C$  such that all points of  $K$  are within  $C$ . As  $S_1$  and  $S_2$  are unbounded, there is a point  $P_1$  of  $S_1$  and a point  $P_2$  of  $S_2$  without  $C$ . Join  $P_1$  and  $P_2$  by an arc lying entirely without  $C$ . By Lemma *A*, this arc must contain a point of  $K$ . But all points of  $K$  are within  $C$ . Hence we are led to a contradiction if we suppose our condition is not necessary.

<sup>\*</sup> Cf. R. L. Moore, *Foundations*, Theorem 39, pp. 153-5.

<sup>†</sup> Cf. R. L. Moore and J. R. Kline, *On the most general closed point-set through which it is possible to pass a simple continuous arc*, *Annals of Mathematics*, vol. XX (1919), pp. 218-23.



*The condition is sufficient.* For suppose  $S_1$  is bounded while  $K$  is unbounded. Since  $S_1$  is bounded, there exists a circle  $C$  enclosing  $S_1$ . Since  $K$  is unbounded, it contains a point  $P$  without  $C$ . The point  $P$  cannot be a limit point of  $S_1$ . But this is contrary to hypothesis.

*Proof of Theorem A.* Two cases may arise:

*Case I.  $K$  is bounded.* Then, by Schoenflies' Theorem and the preceding lemmas, it follows that  $K$  is a simple closed curve.

*Case II.  $K$  is unbounded.* It follows, by Lemma  $F$  that neither  $S_1$  nor  $S_2$  is bounded. Then  $K$  is an open curve. For a proof of this statement see my paper, "The converse of the theorem concerning the division of a plane by an open curve."\*

UNIVERSITY OF PENNSYLVANIA,  
PHILADELPHIA, PA.

---

\* Cf. these Transactions, vol. 18 (1917), pp. 177-184.